

**Comparison of CERES/SSF-MODIS and
CERES/SSF-VIRS Aerosol Products**

X.-P. Tom Zhao and I. Laszlo

NOAA/NESDIS/ORA

27th CERES STM

September 17-19, 2002

- Objective

- Assess aerosol data continuity in the transition from CERES/SSF-VIRS to CERES/SSF-MODIS data sets.

- Approach

- Comparison of results from self-consistency checks.
- Comparison of global daily and monthly mean values.




Data Sets

- **SSF-VIRS only data for March 2001**
- **SSF-MODIS (Terra) Beta1 (FM1,FM2) data for March 2001 (VIRS-like)**




Retrieval Algorithm

- **Input Radiance (different):**
 - VIRS (0.63 μm , 1.61 μm); MODIS (0.67 μm , 1.64 μm)
- **Cloud Screening (same):** Minnis et al. (2001)
- **Sampling (same):** $\gamma > 40^\circ$ & anti-solar side of orbit
- **Aerosol Retrieval (same):** Ignatov & Stowe (2002)
 - dependent two-channel retrieval; Fresnel ($v=1\text{m/s}$) + small Diff. Ref.
 - bi-modal log-normal aerosol distribution
 - LUTs using 6S Code (Vermote et al., 1997)

Probability Distribution of τ




Probability Distribution of α




Global Map of $\Delta\tau$


SSF AOT1 DIFF (MODIS/FM1-MODIS/FM2) for March




SSF AOT1 DIFF (MODIS/FM1-VIROnly) for March



SSF AOT2 DIFF (MODIS/FM1-MODIS/FM2) for March




SSF AOT2 DIFF (MODIS/FM1-VIROnly) for March






Global Map of $\Delta\alpha$

SSF ALPH DIFF (MODIS/FM1-MODIS/FM2) for March



SSF ALPH DIFF (MODIS/FM1-VIROnly) for March



Comparison of Globally Averaged τ_1 , τ_2 , and α for MODIS/SSF and VIRS/SSF in March 2001

	SSF Data	Day1	Day15	Day31	Day 1-10	Day1-31
τ_1	M-FM1	0.118	0.101	0.100	0.110	0.105
	M-FM2	0.097	0.082	0.092	0.098	0.085
	V-Only	0.199	0.135	0.134	0.178	0.139
	V-ED2A	0.176	0.147	0.214	0.138	0.158
τ_2	M-FM1	0.085	0.076	0.070	0.082	0.080
	M-FM2	0.070	0.064	0.064	0.068	0.066
	V-Only	0.088	0.071	0.065	0.085	0.070
	V-ED2A	0.125	0.096	0.115	0.089	0.099
α	M-FM1	0.234	0.151	0.212	0.182	0.152
	M-FM2	0.186	0.039	0.153	0.087	0.062
	V-Only	1.003	0.895	1.032	0.949	0.936
	V-ED2A	0.310	0.442	0.669	0.503	0.509
Note: The time for the VIRS-ED2A data is March 1998.						

Summary

- **AOT and α from FM1 and FM2 of SSF/MODIS are comparable with FM1 values slightly larger.**
- **AOT1 and α of SSF/VIRS are systematically larger than that of SSF/MODIS, especially in the NH for AOT1 (> 30%) and in the SH for α (several factors).**
- **SSF/VIRS AOT2 is comparable to SSF/MODIS AOT2 with slightly larger values of SSF/VIRS in the tropics and slightly smaller values in the subtropics.**

Future Works

- **Apply the same analyses to new version of SSF aerosol products (e.g., TERRA/SSF-ED1)**
- **Explore the possibility of inter-calibration between VIRS and MODIS sensors to find a better solution for the calibration of VIRS channel 2 (1.61 μ m).**

Acknowledgements

**E. Geier, L. Hunt, Langley DAAC
B. Wielicki, B. Barkstrom, CERES
CERES Science Team Members**