

TOA Direct Radiative Effect of Aerosols

Norman G. Loeb

Hampton University/NASA Langley Research Center, VA

Natividad Manalo-Smith

Analytical Services and Materials, Hampton, VA

May 4, 2005, GFDL

Introduction

- Do models and observations provide consistent estimates of the total (natural+anthropogenic) direct radiative effect of aerosols (DREA)?
- **What are the uncertainties in the observations and how can they be reduced?**
- **What is the seasonal and interannual variability in the DREA?**
- What observations are needed to determine the natural and anthropogenic components of the total DREA?

Direct Radiative Effect of Aerosols (Natural+Anthropogenic)


$$\Delta \bar{F}_{SW}^{clr}(\Theta, \Phi) = \bar{F}_{SW}^{na}(\Theta, \Phi) - \bar{F}_{SW}^{clr}(\Theta, \Phi)$$

$\bar{F}_{SW}^{clr}(\Theta, \Phi)$ = clear-sky SW TOA flux

$\bar{F}_{SW}^{na}(\Theta, \Phi)$ = clear-sky SW TOA flux (no aer)

	Spectral Resolution	Spatial Resolution
MODIS	Narrowband	<u>0.5 km</u>
CERES	<u>Broadband</u>	20 km

DIRECT RADIATIVE EFFECT OF AEROSOLS



NARROW-TO-BROADBAND REGRESSIONS

Data:

- CERES Single Scanner Footprint TOA/Surface Fluxes and Clouds (SSF)
- MODIS radiances at 0.644 μm , 0.858 μm , and 1.632 μm
- March 2000 - December 2003


Multi-Channel Regression Fit:

$$\hat{I}_{SW}^{clr} = a_o + \sum_{i=1}^{N_\lambda} a_i I_i^{clr}$$

- I_i cloud-free MODIS radiance in i^{th} channel
- Function of viewing geometry ($\Delta\theta_o=10^\circ$; $\Delta\theta=10^\circ$; $\Delta\phi=20^\circ$)
- a_i 's determined monthly using cloud-free CERES FOVs
- Avoid sunglint


1°x1° Regional Relative Bias and RMS Error in SW Radiance From Narrow-to-Broadband Regression

Relative Bias Error








Avg relative bias error = -0.5%
(=> -0.2 W m⁻² 24-h avg flux)

Relative RMS Error




Avg relative RMS error = 2.75%
(=> 1 W m⁻² 24-h avg flux)

Clear-Sky SW Direct Radiative Effect of Aerosols: Sensitivity to Cloud Mask




Aerosol Direct Radiative Effect (W m^{-2})





Difference in Clear-Sky SW Direct Radiative Effect



Difference in 0.644- μ m Aerosol Optical Depth




0.644- μm Aerosol Optical Depth vs Cloud Fraction (JJA 2000)




MOD04 Aerosol Coverage & 0.644- μm Aerosol Optical Depth in “Overcast” CERES Footprints Identified by CERES Cloud Mask


Aer Covg (%)



$\tau_a(0.644 \mu\text{m})$




Clear-Sky SW Direct Radiative Effect: Comparison with Chou et al. (2002)




Annual Average Clear-Sky SW Direct Radiative Effect:
Comparison with Chou et al. (2002)

	SW Direct Effect of Aerosols ($W m^{-2}$)		
	NOAA-SSF	MOD04	Chou_2002
Global	-3.8	-5.5	-5.4
SH	-3.2	-4.6	-5.1
NH	-4.4	-6.4	-5.7

Stratification by Region





Direct Radiative Effect of Aerosols Over Land


$$\Delta \bar{F}_{SW}^{aer}(\Theta, \Phi) = \bar{F}_{SW}^{na}(\Theta, \Phi) - \bar{F}_{SW}^{clr}(\Theta, \Phi)$$

$\bar{F}_{SW}^{na}(\Theta, \Phi)$ = MODIS Land Albedo + Rayleigh Atmosphere


$\bar{F}_{SW}^{clr}(\Theta, \Phi)$ = CERES clear-sky SW TOA Flux

Global Direct Radiative Effect of Aerosols (CERES+MOD04+MOD43)


JAN, 2001





APR, 2001




JUL, 2001



OCT, 2001



Direct Radiative Effect of Aerosols Over Land (Comparison with Aeronet-Derived Values)



Aeronet-derived DREA provided by Stefan Kinne

Summary

- Global ocean clear-sky SW direct radiative effect of aerosols estimated to be -5.5 W m^{-2} (MOD04) and -3.8 W m^{-2} (NOAA-SSF).
- Large regional and global uncertainty due to cloud mask differences, especially near desert regions.
-> CALIPSO should help with this.
- The DREA has pronounced seasonal cycle in the Northern Hemisphere and large year-to-year fluctuations between 30° - 60° N.
- No systematic trend in deseasonalized anomalies of the DREA is observed over the 46-month time series considered