

Heating of Earth's Climate System Continues Despite Slowing of Surface Warming in Past Decade

Norman G. Loeb¹, John M. Lyman², Gregory C. Johnson², Richard P. Allan³, David R. Doelling¹, Takmeng Wong¹, Brian J. Soden⁴, Fred Rose⁵, and Graeme L. Stephens⁵

¹NASA LaRC

²NOAA

³U. Reading

⁴U. Miami

⁵SSAI


⁶JPL

Global Surface Temperature Anomalies

Source: HadCRUT3

1981–2010: 0.016 °C/yr


2001–2010: 0.000 °C/yr



Source: GISSTEMP

1981–2010: 0.021 °C/yr


2001–2010: 0.016 °C/yr



Source: ERA-Interim

1981–2010: 0.014 °C/yr

2001–2010: 0.005 °C/yr




What do the new generation of observations (satellite, ocean) tell us about the associated energy flow during the past decade?


Tracking Earth's Energy

- Measuring net incoming and outgoing radiation at TOA indicates how much energy remains in the Earth system.
- Most of this excess energy (~90%) is absorbed in the ocean; rest goes into melting sea and land ice and heating the land surface and atmosphere.
- Fluctuations in Earth's TOA net energy imbalance and the ocean heating rate should be in phase with one another.
- Sources of short-term variability: ENSO-related changes in atmospheric and oceanic circulations, clouds, aerosols (including volcanic eruptions), surface properties.
- Longer-term variability: Anthropogenic forcing due to greenhouse gases and aerosols, natural forcing by aerosols and solar radiation, and feedbacks involving water vapor, temperature, clouds, and the surface.
- Earth's temperature response depends upon the thermal inertia of the climate system, governed by the ocean (large heat capacity).


The Sun, Stratospheric Aerosols, and Lower-Stratospheric Water Vapor



Total solar irradiance changes: -0.5 Wm^{-2}
 \Rightarrow reduction in absorbed solar radiation
of 0.1 Wm^{-2} (Trenberth, 2009)




Lower stratospheric water vapor
changes: -0.4 ppmv
 $\Rightarrow 0.04^\circ\text{C}$ reduction in warming
Solomon et al. (2010)




Stratospheric aerosol
changes: $+7\%$ per yr
 $\Rightarrow 0.07^\circ\text{C}$ reduction in
warming
Solomon et al. (2011)


CERES SW, LW and Net TOA Radiation & ENSO



AIRS Temperature and Water Vapor Mixing Ratio Anomalies (30°S-30°N)



Temperature
Anomaly (K)



Water Vapor Mixing Ratio
Anomaly (g/kg)

Loeb et al., 2011 (Surv. Geophys.)






Figure 4 Monthly anomalies in (a) CERES Terra SW TOA flux from SSF1deg-lite_Ed2.5 and SeaWiFS PAR scaled by a factor of -6.09 (corresponding to the slope of the regression line fit relating CERES SW TOA flux and SeaWiFS PAR anomalies) over ocean for $30^{\circ}S-30^{\circ}N$ from March 2000 to December 2009, (b) CERES Terra SW TOA flux and MODIS cloud fraction for $30^{\circ}S-30^{\circ}N$ between March 2000 and February 2010, and (c) global LW TOA flux from CERES Terra, CERES Aqua and AIRS Aqua for September 2002-February 2010.




Apparent inconsistencies after 2004 between short-term variations in ocean heating rate from in situ ocean heat content data and net TOA flux from CERES cast doubt on our ability to account for flows of energy in the climate system, and the lack of closure has given rise to the idea of “missing energy” in the climate system.

Global Average Upper Ocean Heating Rates from 3 Different Teams




Comparison of net TOA flux and Upper Ocean Heating Rates




CERES TOA Flux and MODIS Cloud Property Anomalies (Feb 2008)


CERES SW TOA Flux Anomaly (Wm^{-2})




CERES LW TOA Flux Anomaly (Wm^{-2})



MODIS Cloud Fraction Anomaly (%)



MODIS Cloud Top Pressure Anomaly (hPa)



Diagnosing CERES Tropical Mean LW TOA Flux Variability

Consider the time series of computed monthly mean LW TOA flux:

$$F^{LW}(T(z), w(z), T_s, P_c, f, \tau_c)$$

Compute contribution to LW TOA flux variability from each variable:

$$\delta F^{LW}(T(z)) = F^{LW}(T(z), w(z), T_s, P_c, f, \tau_c) - F^{LW}(\bar{T}(z), w(z), T_s, P_c, f, \tau_c)$$

$$\delta F^{LW}(w(z)) = F^{LW}(T(z), w(z), T_s, P_c, f, \tau_c) - F^{LW}(T(z), \bar{w}(z), T_s, P_c, f, \tau_c)$$


$$\delta F^{LW}(T_s) = F^{LW}(T(z), w(z), T_s, P_c, f, \tau_c) - F^{LW}(T(z), w(z), \bar{T}_s, P_c, f, \tau_c)$$

$$\delta F^{LW}(P_c) = F^{LW}(T(z), w(z), T_s, P_c, f, \tau_c) - F^{LW}(T(z), w(z), T_s, \bar{P}_c, f, \tau_c)$$

$$\delta F^{LW}(f) = F^{LW}(T(z), w(z), T_s, P_c, f, \tau_c) - F^{LW}(T(z), w(z), T_s, P_c, \bar{f}, \tau_c)$$

$$\delta F^{LW}(\tau_c) = F^{LW}(T(z), w(z), T_s, P_c, f, \tau_c) - F^{LW}(T(z), w(z), T_s, P_c, f, \bar{\tau}_c)$$

Contributions to LW TOA Flux by Surface, Atmos & Clouds (30S-30N)



- Cloud and atmos contributions in phase during strong ENSO.
- Cloud fraction decrease & mean cloud-top pressure increase during El Nino (-ve LW fdbck; +ve SW fdbk)

Conclusions

- We calculate that during the past decade Earth has been accumulating energy at the rate $0.58 \pm 0.43 \text{ Wm}^{-2}$, suggesting that while warming of Earth's surface has slowed during the 2000s, energy continues to accumulate in the subsurface ocean.
- The apparent decline in ocean heating rate after 2004 noted in other studies is not statistically robust.
- Our results do not support the claim that there is missing energy in the system.
- Synergistic use of satellite and in-situ ocean heat content anomaly measurements provide critical data for quantifying short and longer-term changes in the Earth's net TOA radiation imbalance.
- New cloud data from satellite (MODIS, CALIPSO) key to unraveling role of clouds in a changing climate.